

# MAXIM<sup>®</sup>

## Lemon Detergent Disinfectant



DS 470

### DESCRIPTION

When used as directed, this disinfectant is formulated to disinfect inanimate hard surfaces such as walls, floors, sink tops, toilet bowls, tables, chairs, telephones, and bed frames. For larger areas such as operating rooms, patient care facilities and restrooms, this product is designed to provide both general cleaning and disinfecting. Effective against HIV-1 (AIDS), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), MRSA, and Trichophyton mentagrophytes (athlete's foot fungus).

### DIRECTIONS

Read entire container label before using this product.  
This product is not for use on Medical device surfaces.

**HOSPITALS, DENTAL OFFICES, NURSING HOMES AND OTHER HEALTH CARE INSTITUTIONS:** Add 2 ounces of this product to one gallon of water. Treated surfaces must remain wet for 10 minutes. Apply this product with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surfaces must be sprayed until thoroughly wetted. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray. NOTE: With spray applications, cover or remove all food products. Prepare a fresh solution at least daily or when use solution becomes visibly dirty. For heavily soiled areas, a preliminary cleaning is required.

When used as directed this disinfectant exhibits effective disinfectant activity against the following: *Campylobacter jejuni*, *Corynebacterium ammoniagenes*, *Enterobacter aerogenes*, *Enterococcus faecalis*, *Escherichia coli*, *Listeria monocytogenes*, *Pseudomonas aeruginosa*, *Salmonella enterica*, *Salmonella typhi*, *Shigella dysenteriae*, *Staphylococcus aureus*, *Staphylococcus aureus*—Methicillin resistant, *Streptococcus salivarius*, *Aspergillus niger*, *Trichophyton mentagrophytes*—athlete's foot fungus, Avian Influenza A H5N1 virus, Hepatitis B Virus, Hepatitis C Virus, Herpes Simplex Type1, Human Immunodeficiency Virus Type I—HIV-1, Influenza A2/Japan virus, Norovirus.

### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

### APPLICATIONS

Cleaning, disinfecting, and deodorizing in:

- Hospitals
- Nursing homes
- Schools

### BENEFITS

- Use on all washable surfaces
- Cleans and disinfects
- Excellent deodorizer
- Highly dilutable

### SPECIFICATIONS

Color..... Yellow  
Fragrance.....Lemon  
pH Range..... 11.5-12.0  
Dilution .....2 oz./gal.  
Density .....8.65  
EPA Registered..... Yes

### PACKAGING

ITEM # .....SIZE  
047000-41 .....4/1 gal.  
047000-05 .....5 gal.  
047000-55 .....55 gal.

**MIDLAB**  
INCORPORATED

232 Industrial Park Rd  
Sweetwater, TN 37874  
1.800.467.6294  
midlab.com